

High Resolution Time Series Observations of Bio-Optical and Physical Variability in the Arabian Sea

Tommy D. Dickey
Ocean Physics Laboratory
University of California at Santa Barbara
6487 Calle Real, Suite A
Goleta, CA 93117

phone: (805) 893-7354 fax: (805) 967-5704 email tommy@icess.ucsb.edu

Award # N00014-96-1-0505

<http://www.icess.ucsb.edu/opl/arabian.html>

LONG-TERM GOALS

An interdisciplinary mooring field study was conducted in the Arabian Sea (October 1994 - November 1995) as part of the ONR Forced Upper Ocean Dynamics (FUOD) Program activities in the region. The Arabian Sea experiment allowed us to study bio-optical and physical processes and their interactions under monsoonal and inter-monsoonal conditions. This region is attractive because of its regular and intense atmospheric forcing which causes extremely strong responses in both the physics and biology of the upper layer. Thus, dynamical ranges in measured properties are great, enabling us to apply and test time-dependent interdisciplinary models relevant to bio-optical properties and carbon fluxes.

SCIENTIFIC OBJECTIVES

Our work addresses several scientific questions, allows testing of specific hypotheses, and provides complementary data for collaborating scientists. Key questions include:

1. What are the dominant scales of coupling between atmospheric forcing and physical/bio-optical responses (e.g., mixed layer and stratification evolution, phytoplankton blooms and busts)?
2. What are the roles of episodic wind events, monsoons, and seasonal insolation with respect to biological/physical processes and their couplings?
3. To what degree can physical/bio-optical/biological dynamics be explained using local (one-dimensional) models? What are the responses of the mixed and euphotic layers to advective features such as filaments and eddies?
4. How can horizontal information (e.g., satellite surface temperature, topography, and color data, ship-based underway and tow-yo data) be synthesized with mooring data?
5. Do phytoplankton blooms significantly affect water column heating, stratification, and horizontal currents?

APPROACH

The ONR Forced Upper Ocean Dynamics (FUOD) Program utilized an array of 5 deep ocean moorings placed beneath the axis of the Findlater Jet to study linkages between large-scale patterns of wind speed and direction, ocean circulation, and biological production in near surface waters. The central (Woods Hole Oceanographic Institution, WHOI) mooring was located at 15° 30'N, 61° 30'E, the center of a 50 km square with other moorings placed at the corners. The observations of

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE High Resolution Time Series Observations of Bio-Optical and Physical Variability in the Arabian Sea				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California at Santa Barbara, Ocean Physics Laboratory, 6487 Calle Real, Suite A, Goleta, CA, 93117				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

primary interest for the present work were made from instruments deployed from the central (WHOI) mooring during two separate deployments (October 15, 1994-April 20, 1994 and April 22, 1995-October 20, 1995). Multi-variable moored systems (MVMS) were deployed by our group at 35 and 80m. The MVMS utilizes a VMCM to measure currents and temperature with a pressure case mounted thermistor. A variety of other sensors are used to measure photosynthetic available radiation (PAR), natural fluorescence or upwelling radiance at 683nm, stimulated fluorescence, beam attenuation coefficient, and dissolved oxygen. Conductivity was also measured at 35m. All data were sampled at 3.75min intervals. Two other MVMSs were deployed by LDEO at depths of 10 and 65m. The sensor suite was similar to that of the UCSB MVMSs.

WORK COMPLETED

Our MVMS interdisciplinary systems with sampling intervals of a few minutes were placed on a mooring beneath the axis of the Findlater Jet in October 1994 for 6 months and again in April of 1995. Remote sensing data, ship ADCP data and ship tow-yo data have been used to assist in the interpretation of this highly variable environment. The data return from the central mooring was very good considering such a harsh environment to operate underwater electronics. Three data reports, a M.S. thesis, and five papers are in press in special volumes of Deep-Sea Research (please see references below).

RESULTS

The FUOD mooring instrumented with optical and physical sensors within the upper 300 m was deployed for two consecutive 6-month periods (October 15, 1994 through October 20, 1995; sampling intervals of a few minutes) in the central Arabian Sea (15° 30'N, 61° 30'E). Both the northeast and southwest monsoons were observed. During the NE monsoon, wind speeds averaged 6 m/s and reached up to 15 m/s during the SW monsoon. Inter-monsoons were characterized by weak and variable winds. Shortwave radiation and photosynthetically available radiation (PAR) displayed biannual cycles, peaking during the inter-monsoon periods. The maximum winter mixed layer depth (MLD1.0°C~110 m) was deeper than the summer mixed layer (MLD1.0°C~80 m), primarily because of surface cooling and convection. A biannual cycle in chlorophyll was evident with greater values occurring during each monsoon and into the inter-monsoon periods. High chlorophyll values associated with cool mesoscale features were also apparent during each monsoon. These mesoscale features and others have been documented using remotely sensed sea-surface height anomaly maps. Time series of the 1% light level depth, h1%, tracked the depth-integrated chlorophyll. In general, h1% was deeper than MLD1.0°C during the latter half of the spring inter-monsoon (low chlorophyll periods) and shallower than h1% during the latter portions of the monsoons (high chlorophyll periods). The highest mixed layer radiant heating rates occurred during the inter-monsoon periods with peak values greater than 0.15 °C d⁻¹ for MLD1.0°C. These values are consistent with those previously suggested for the central Arabian Sea. Our results indicate that biological variability is important for the seasonal variability of the upper ocean heat budget of the

IMPACT/APPLICATION

Improved understanding of temporal variations of the physical and optical properties of the upper ocean in the Arabian Sea has resulted from our advanced sampling systems and analyses. Predictive modeling of interdisciplinary processes is being made possible through our research efforts. Our work is also important for relating upper ocean particle fluxes with those at depth.

TRANSITIONS

Results from the present work will be valuable for developing models of physical and optical variability in complex regions where intense atmospheric forcing occurs, differing water masses merge, and mesoscale features are ubiquitous and energetic.

RELATED PROJECTS

One of the great strengths of the Arabian Sea program was the large number of collaborating scientists whose data sets complement our own and facilitate the interpretation of our observations. Complementary shipboard data have proven valuable. In particular, vertical profiling and tow-yoing (SeaSoar) of bio-optical sensors and physical and satellite data indicating surface features in ocean temperature and currents have been useful for our interpretations. In addition, a sediment trap mooring located near our mooring array provided information concerning flux of particulate matter to depth. The collective data sets are critical for bio-optical and physical modelers.

REFERENCES

Dickey, T., J. Marra, D. Sigurdson, R. Weller, C. Kinkade, S. Zedler, J. Wiggert, and C. Langdon, 1998, Seasonal variability of bio-optical and physical properties in the Arabian Sea, October 1994 - October 1995, in press, Deep-Sea Research.

Kinkade, C.S., J. Marra, T. Dickey, C. Langdon, D. Sigurdson, and R. Weller, 1998, Diel bio-optical variability in the Arabian Sea as observed from moored sensors, in press, Deep-Sea Research.

Kinkade, C.S., J. Marra, T. Dickey, and R. Weller, 1998, An intercomparison of moored optical sensor data and an annual cycle of phytoplankton biomass in the Arabian Sea, 1994-1995, in press, Deep-Sea Research.

Marra, J. T. Dickey, C. Ho, C. Kinkade, R. Weller, and R. Barber, 1998, Variability in primary production as observed from moored sensors in the central Arabian Sea in 1995, in press, Deep-Sea Research.

Rudnick, D.L., R.A. Weller, C.C. Eriksen, T.D. Dickey, J. Marra, and C. Langdon, 1997, Moored instruments weather Arabian Sea monsoons, yield data EOS, 78, 117, 120-121.

Sigurdson, D.E., 1996, Analysis of physical and bio-optical variability in the Arabian Sea during the northeast monsoon of 1994, M.S. thesis, University of Southern California.

Sigurdson, D.E., Arabian Sea TOPEX data report: September 16, 1994 - September 18, 1995, University of California at Santa Barbara Ocean Physics Laboratory Report OPL-3-96.

Sigurdson, D.E., T.D. Dickey, and D.V. Manov, 1995, Arabian Sea Mooring Data Report: Deployment #1, October 15, 1994 - April 20, 1995, University of California at Santa Barbara Ocean Physics Laboratory Report OPL-2-95, 26pp.

Sigurdson, D.E., T.D. Dickey, and D.V. Manov, 1996, Arabian Sea Mooring Data Report: Deployment #2, April 22, 1995 - October 20, 1995, University of California at Santa Barbara Ocean Physics Laboratory Report OPL-2-96, 26pp.

Wiggert, J., B. Jones, T. Dickey, K. Brink, R. Weller, J. Marra, and L. Codispoti, 1998, The Northeast Monsoon's impact on mixing, phytoplankton biomass, and nutrient cycling in the Arabian Sea, in press, Deep-Sea Research.